

SIMPLE BOUNDS FOR SOLUTIONS OF MONOTONE
COMPLEMENTARITY PROBLEMS AND CONVEX PROGRAMS

by

O. L. Mangasarian & L. McLinden

Computer Sciences Technical Report #534

February 1984

UNIVERSITY OF WISCONSIN-MADISON
COMPUTER SCIENCES DEPARTMENT

SIMPLE BOUNDS FOR SOLUTIONS OF MONOTONE
COMPLEMENTARITY PROBLEMS AND CONVEX PROGRAMS

O. L. Mangasarian, L. McLinden

Technical Report #534

February 1984

Revised August 1984

ABSTRACT

For a solvable monotone complementarity problem we show that each feasible point which is not a solution of the problem provides simple numerical bounds for some or all components of all solution vectors. Consequently for a solvable differentiable convex program each primal-dual feasible point which is not optimal provides simple numerical bounds for some or all components of all primal-dual solution vectors. We also give an existence result and simple bounds for solutions of monotone complementarity problems satisfying a new, distributed constraint qualification. This result carries over to a simple existence and boundedness result for differentiable convex programs satisfying a similar constraint qualification.

AMS (MOS) Classification: 90C30, 90C25

Key Words: Nonlinear programming, complementarity, monotonicity,
convexity, bounds

SIMPLE BOUNDS FOR SOLUTIONS OF MONOTONE
COMPLEMENTARITY PROBLEMS AND CONVEX PROGRAMS

O. L. Mangasarian, L. McLinden

1. The Monotone Complementarity Problem

This work is based on an extremely simple, but apparently unnoticed, property of the monotone complementarity problem [2,5,8,11,12] of finding a (z,w) in the $2k$ -dimensional Euclidean space \mathbb{R}^{2k} such that

$$(1.1) \quad w = F(z) \geq 0, \quad z \geq 0, \quad zw = 0$$

Here $F: D \rightarrow \mathbb{R}^k$ is a monotone function on D where $\mathbb{R}_+^k \subset D \subset \mathbb{R}^k$, that is

$$(z^2 - z^1)(F(z^2) - F(z^1)) \geq 0 \quad \text{for all } z^1, z^2 \in D$$

The property is the following:

1.1 Theorem Let (z,w) be some feasible point of a solvable monotone complementarity problem (1.1), that is $w = F(z) \geq 0, z \geq 0$. Any solution (\bar{z}, \bar{w}) of (1.1) is bounded as follows:

$$(a) \quad \|\bar{z}_I\|_1 := \sum_{i \in I} \bar{z}_i \leq zw / \min_{i \in I} w_i := zw / \min_{i \in I} w_i \quad (I \neq \emptyset)$$

$$(b) \quad \|\bar{w}_J\|_1 \leq zw / \min_{i \in J} z_i \quad (J \neq \emptyset)$$

$$(c) \quad \|\bar{z}_I, \bar{w}_J\|_1 \leq zw / \min \{z_{i \in J}, w_{i \in I}\} \quad (I \cup J \neq \emptyset)$$

where $I = \{i | w_i > 0\}$ and $J = \{i | z_i > 0\}$.

Proof For any solution (\bar{z}, \bar{w}) of (1.1) we have by the monotonicity of F and $\bar{z}\bar{w} = 0$ that

$$zw \geq \bar{z}w + z\bar{w}$$

Hence by the nonnegativity of (z, w) and (\bar{z}, \bar{w}) we have

- (a) $zw \geq \bar{z}_I w_I \geq \|\bar{z}_I\|_1 \min_{i \in I} w_i$
- (b) $zw \geq z_J \bar{w}_J \geq \|\bar{w}_J\|_1 \min_{i \in J} z_i$
- (c) $zw \geq \bar{z}_I w_I + z_J \bar{w}_J \geq \|\bar{z}_I, \bar{w}_J\|_1 \cdot \min \{z_{i \in J}, w_{i \in I}\} \quad \square$

Theorem 1.1 is a partial extension to the monotone complementarity problem of a corresponding result, Theorem 2.2 of [7], for the positive semidefinite linear complementarity problem. Note that, unlike the linear case, feasibility for the nonlinear monotone complementarity problem does not imply solvability as shown by the simple example of [10].

Theorem 1.1 shows that any feasible point (z, w) of a solvable monotone complementarity problem (1.1) which is not a solution of the problem (so that both I and J are nonempty) provides some information about the magnitude of the solution set. In certain cases, such as when $w > 0$, we get a bound on all components of all solution vectors \bar{z} .

With the bounds given by Theorem 1.1 it is possible to obtain bounds for optimal solutions and multipliers of solvable differentiable convex programs once they are cast as monotone complementarity problems. (See Section 2.) But before doing that we show how the bounds of Theorem 1.1 can be extended to approximate solutions of monotone complementarity problems which may not even be solvable. Let

$$(1.2) \quad \alpha := \inf \{zw \mid w = F(z) \geq 0, z \geq 0\} \geq 0,$$

and for $\epsilon \geq 0$ let $(\bar{z}(\epsilon), \bar{w}(\epsilon))$ be an ϵ -solution of the optimization problem of (1.2), that is

$$(1.3) \quad \bar{w}(\epsilon) = F(\bar{z}(\epsilon)) \geq 0, \bar{z}(\epsilon) \geq 0, \alpha + \epsilon \geq \bar{z}(\epsilon)\bar{w}(\epsilon) \geq \alpha$$

Note that for any $\epsilon > 0$, an ϵ -solution always exists provided problem (1.2) has at least one feasible point. A 0-solution exists provided the infimum of (1.2) is attained, that is the infimum is a minimum. Furthermore if $\alpha = 0$, then an ϵ -solution of (1.2) is an "approximate" solution of the complementarity problem (1.1) which is an exact solution if $\epsilon = 0$. With these concepts in mind it follows from (1.2), (1.3) and the monotonicity of F that, for any feasible (z, w) and $\epsilon \geq 0$,

$$(1.4) \quad 2zw + \epsilon \geq zw + \alpha + \epsilon \geq zw + \bar{z}(\epsilon)\bar{w}(\epsilon) \geq z\bar{w}(\epsilon) + \bar{z}(\epsilon)w$$

Consequently a generalization of Theorem 1.1 is possible if, instead of one feasible point (z, w) , we consider p the feasible points (z^j, w^j) , $j=1, 2, \dots, p$, of the optimization problem of (1.2) and corresponding weights $\lambda^j \geq 0$, $j=1, \dots, p$, such that $\sum_{j=1}^p \lambda^j = 1$. Then by (1.4) we have that

$$(1.5) \quad 2 \sum_{j=1}^p \lambda^j z^j w^j + \epsilon \geq \sum_{j=1}^p \lambda^j z^j w^j + \alpha + \epsilon \geq \sum_{j=1}^p \lambda^j z^j w^j + \bar{z}(\epsilon)\bar{w}(\epsilon) \geq \sum_{j=1}^p \lambda^j z^j \bar{w}(\epsilon) + \sum_{j=1}^p \lambda^j \bar{z}(\epsilon) w^j$$

Then, arguing as in Theorem 1.1 we obtain the following bounds.

1.2 Theorem Let F be monotone on R_+^k and let (z^j, w^j) , $j=1, 2, \dots, p$, be feasible points of the optimization problem of (1.2), that is, $w^j = F(z^j) \geq 0$,

$z^j \geq 0, j=1,2,\dots,p$. Let $\lambda^j \geq 0, j=1,2,\dots,p, \sum_{j=1}^p \lambda^j = 1$ and let $\epsilon \geq 0$. Any ϵ -solution $(\bar{z}(\epsilon), \bar{w}(\epsilon))$ of (1.2) defined by (1.3) is bounded as follows:

$$(a) \quad \|\bar{z}_I(\epsilon)\|_1 \leq \left(\sum_{j=1}^p \lambda^j z^j w^j + \alpha + \epsilon \right) / \min_{i \in I} \hat{w}_i \leq \left(2 \sum_{j=1}^p \lambda^j z^j w^j + \epsilon \right) / \min_{i \in I} \hat{w}_i$$

$$(b) \quad \|\bar{w}_J(\epsilon)\|_1 \leq \left(\sum_{j=1}^p \lambda^j z^j w^j + \alpha + \epsilon \right) / \min_{i \in J} \hat{z}_i \leq \left(2 \sum_{j=1}^p \lambda^j z^j w^j + \epsilon \right) / \min_{i \in J} \hat{z}_i$$

$$(c) \quad \|\bar{z}_I(\epsilon), \bar{w}_J(\epsilon)\|_1 \leq \left(\sum_{j=1}^p \lambda^j z^j w^j + \alpha + \epsilon \right) / \min\{\hat{z}_{i \in J}, \hat{w}_{i \in I}\} \leq \left(2 \sum_{j=1}^p \lambda^j z^j w^j + \epsilon \right) / \min\{\hat{z}_{i \in J}, \hat{w}_{i \in I}\}$$

where $I = \{i | \hat{w}_i > 0\}$, $J = \{i | \hat{z}_i > 0\}$, $\hat{z} := \sum_{j=1}^p \lambda^j z^j$ and $\hat{w} := \sum_{j=1}^p \lambda^j w^j$.

We note that the first inequality of each of (a), (b) and (c) of Theorem 1.2 remains valid even if we do not require that $w^j \geq 0$ and $z^j \geq 0$, but merely that $z^j \in D$, where $R_+^k \subset D \subset R^k$, F is monotone on D and $\hat{z} \geq 0$ and $\hat{w} \geq 0$. This remark will be employed in Theorem 1.3.

The bounds established by Theorem 1.2(a) provide motivation for the proof of the following existence result, which employs a new "distributed" constraint qualification.

1.3 Theorem (Existence and boundedness of solutions of monotone complementarity problems under a distributed constraint qualification) Let

$F: D \rightarrow R^k$ be monotone and continuous on D such that $R_+^k \subset D \subset R^k$, let

$z^j \in D, w^j = F(z^j) \in R^k, j=1,2,\dots,p$, be such that $\hat{z} := \sum_{j=1}^p \lambda^j z^j \geq 0$,

$\hat{w} := \sum_{j=1}^p \lambda^j w^j > 0$ for some $\lambda^j \geq 0, j=1,2,\dots,p, \sum_{j=1}^p \lambda^j = 1$. Then the complementarity problem (1.1) is solvable. Any solution (\bar{z}, \bar{w}) is bounded as follows:

$$(1.6) \quad \|\bar{z}\|_1 \leq \left(\sum_{j=1}^p \lambda^j z^j w^j \right) / \min_{1 \leq i \leq k} \hat{w}_i$$

Proof The bound (1.6) follows from Theorem 1.2(a) with $\alpha = \epsilon = 0$ and the remark following it, once we have established the existence of a solution to the complementarity problem (1.1), which we proceed to do now by means of the Brouwer fixed point theorem [1,14]. Let

$$C := \{z \mid z \geq 0, \hat{w}z \leq \hat{w}\hat{z} + \gamma\},$$

where

$$(1.7) \quad \gamma > \max \{1, -\hat{w}\hat{z} + \sum_{j=1}^p \lambda^j z^j w^j\} \geq 1$$

The set C is nonempty, compact and convex and the single-valued mapping [4] defined by the 2-norm projection of $z - F(z)$ on C :

$$z \rightarrow \operatorname{argmin}_{y \in C} \|y - z + F(z)\|_2$$

defines a continuous function from C into itself. Hence by Brouwer's theorem this function must have a fixed point $\bar{z} \in C$. Such a point satisfies the minimum principle optimality condition [6]

$$(1.8) \quad \bar{z} \in C, F(\bar{z})(y - \bar{z}) \geq 0 \quad \forall y \in C$$

If $\hat{w}\bar{z} < \hat{w}\hat{z} + \gamma$ then \bar{z} solves (1.1). Indeed $\bar{z} + \delta e_i$, $i=1,2,\dots,k$, is in C for δ sufficiently small and positive and e_i the i th unit coordinate vector, and hence by (1.8) it follows that $F(\bar{z}) \geq 0$, $\bar{z} \geq 0$, and $\bar{z}F(\bar{z}) \leq 0$ by taking $y = 0$ in (1.8). We now show that the case

$$(1.9) \quad \hat{w}\bar{z} = \hat{w}\hat{z} + \gamma$$

cannot occur. For if it did, then from the monotonicity of F we have

$$\bar{z}F(\bar{z}) \geq -z^j w^j + z^j \bar{w} + \bar{z} w^j, \quad j=1,2,\dots,p$$

where $\bar{w} := F(\bar{z})$. Multiplying by λ^j and summing over j gives

$$\begin{aligned}\bar{z}F(\bar{z}) &\geq \sum_{j=1}^p -\lambda^j z^j w^j + \hat{z}\bar{w} + \bar{z}\hat{w} \\ &> \hat{z}\bar{w} = \hat{z}F(\bar{z}) \quad (\text{By (1.9) and (1.7)})\end{aligned}$$

Hence $F(\bar{z})(\hat{z} - \bar{z}) < 0$ which contradicts (1.8). So (1.9) cannot occur and \bar{z} solves (1.1). \square

We note that the existence part of the above theorem for the ordinary constraint qualification, that is $p = 1$, was obtained by Moré [12, Theorem 3.2] and by one of the authors in [8, Theorem 1] for the case of multivalued monotone mappings.

It is interesting to note that the complementarity problem of Megiddo [10] which has no solution, does not satisfy the distributed constraint qualification of Theorem 1.4 and hence indicates the sharpness of that condition. On the other hand Theorem 1.2(a) can be used to give an exact upper bound on the bounded component of the solution of problem (1.2) for Megiddo's example.

We also note the distributed constraint qualification of Theorem 1.3 is implied by the ordinary constraint qualification if we take $p = 1$. The converse is true if $D = \mathbb{R}_+^k$ and F is concave on \mathbb{R}_+^k . However F is not concave in general, and in fact is merely monotone when it is derived from a differentiable convex program. (See Section 2.) However for the general case of a monotone F and $D = \mathbb{R}_+^k$, it can be shown [9, Theorem 4] that the two constraint qualifications are equivalent. Nevertheless the distributed qualification may be easier to verify.

2. Bounds for Solutions of Convex Programs

We consider now the solvable differentiable convex program

$$(2.1) \quad \min_x f(x) \quad \text{s.t. } y = -g(x) \geq 0, x \geq 0$$

where $f: \mathbb{R}^n \rightarrow \mathbb{R}$, $g: \mathbb{R}^n \rightarrow \mathbb{R}^m$ are convex and differentiable functions, together with its dual [6]

$$(2.2) \quad \max_{x,u} L(x,u) - x \nabla_x L(x,u) \quad \text{s.t. } v = \nabla_x L(x,u) \geq 0, u \geq 0$$

where $L(x,u)$ is the standard Lagrangian

$$L(x,u) = f(x) + ug(x)$$

and ∇_x denotes the gradient vector with respect to x . We note that the Karush-Kuhn-Tucker conditions

$$(2.3) \quad \begin{aligned} v = \nabla_x L(x,u) &= \nabla f(x) + u \nabla g(x) \geq 0, x \geq 0, xv = 0 \\ y = -\nabla_u L(x,u) &= -g(x) \geq 0, u \geq 0, uy = 0 \end{aligned}$$

hold if and only if (x,y,u,v) solves the dual programs (2.1)-(2.2) with equal extrema [6]. If the constraints of (2.1) satisfy the Slater constraint qualification, that is $g(x) < 0$ for some $x \geq 0$, then for each solution of (2.1) the Karush-Kuhn-Tucker conditions (2.3) are satisfiable [6]. If we make the definitions

$$(2.4) \quad z := \begin{pmatrix} x \\ u \end{pmatrix}, \quad w := \begin{pmatrix} v \\ y \end{pmatrix}, \quad F(z) := \begin{pmatrix} \nabla_x L(x,u) \\ -\nabla_u L(x,u) \end{pmatrix}$$

then the Karush-Kuhn-Tucker conditions take on the equivalent complementarity problem formulation [2]

$$(2.5) \quad w = F(z) \geq 0, \quad z \geq 0, \quad zw = 0$$

Note that the monotonicity of the "twisted" derivative involved in the definition of $F(z)$ has also been used in [3,13,5,8]. For completeness we include a proof of this fact.

2.1 Lemma Let f and g be differentiable and convex on R^n and let $F(z)$ be defined as in (2.4). Then $F(z)$ is monotone and continuous for all $z \in R^n \times R_+^m$.

Proof By the convexity of g and $\bar{u} \geq 0, u \geq 0$ we have that

$$\begin{aligned} \bar{u}(g(x) - g(\bar{x})) &\geq \bar{u}\nabla g(\bar{x})(x - \bar{x}) \\ -u(-g(\bar{x}) + g(x)) &\geq -u\nabla g(x)(-\bar{x} + x) \end{aligned}$$

Addition of these two inequalities gives

$$(2.6) \quad -(u - \bar{u})(g(x) - g(\bar{x})) \geq (\bar{u}\nabla g(\bar{x}) - u\nabla g(x))(x - \bar{x})$$

Hence

$$\begin{aligned} (z - \bar{z})(F(z) - F(\bar{z})) &= (x - \bar{x} \quad u - \bar{u}) \begin{pmatrix} \nabla_x L(x, u) - \nabla_x L(\bar{x}, \bar{u}) \\ - (g(x) - g(\bar{x})) \end{pmatrix} \\ &\geq (x - \bar{x})(\nabla f(x) - \nabla f(\bar{x})) \quad (\text{By (2.6)}) \\ &\geq 0 \quad (\text{By convexity of } f) \end{aligned}$$

The continuity of F follows from the fact that a differentiable convex function on R^n is continuously differentiable. \square

We can now apply Theorem 1.1 to the monotone function $F(z)$ of (2.4) to obtain bounds for optimal solutions and multipliers of (2.1).

2.2 Theorem Let f and g be differentiable and convex on R^n . Each primal-dual feasible point of (2.1)-(2.2), that is (x, y, u, v) satisfying

$$y = -g(x) \geq 0, x \geq 0, v = \nabla_x L(x, u) \geq 0, u \geq 0,$$

bounds any point $(\bar{x}, \bar{y}, \bar{u}, \bar{v})$ which solves the primal-dual programs (2.1)-(2.2) with equal extrema, or equivalently, which satisfies the Karush-Kuhn-Tucker conditions (2.3) for (2.1) as follows:

- (a) $\sum_{i \in I_1} \bar{x}_i =: \|\bar{x}_{I_1}\|_1 \leq (xv + uy) / \min_{i \in I_1} v_i$
- (b) $\|\bar{y}_{J_2}\|_1 \leq (xv + uy) / \min_{i \in J_2} u_i$
- (c) $\|\bar{u}_{I_2}\|_1 \leq (xv + uy) / \min_{i \in I_2} y_i$
- (d) $\|\bar{v}_{J_1}\|_1 \leq (xv + uy) / \min_{i \in J_1} x_i$

where

$$I_1 = \{i | v_i > 0\}, J_2 = \{i | u_i > 0\}, I_2 = \{i | y_i > 0\}, J_1 = \{i | x_i > 0\}$$

Proof Immediate from Theorem 1.1, Lemma 2.1 and definition (2.4). \square

Theorem 2.2 is a partial extension of Theorem 3.1 of [7] where bounds for solutions of linear programs were given.

All the other theorems of Section 1 apply in a straightforward manner to the convex program (2.1) via the complementarity formulation (2.4)-(2.5). We state below the counterpart of Theorem 1.3 for the convex program (2.1).

2.3 Theorem (Existence and boundedness of solutions of differentiable convex programs under a distributed constraint qualification)

Let f and g be differentiable and convex on R^n , let

$$y^j = -g(x^j) \in R^m, x^j \in R^n, v^j = \nabla_x L(x^j, u^j) \in R^n, u^j \geq 0, j=1,2,\dots,p$$

be such that for some $\lambda^j \geq 0, j=1,2,\dots,p, \sum_{j=1}^p \lambda_j = 1$:

$$\hat{x} := \sum_{j=1}^p \lambda^j x^j \geq 0, \hat{y} := \sum_{j=1}^p \lambda^j y^j > 0, \hat{v} := \sum_{j=1}^p \lambda^j v^j > 0$$

Then there exists $(\bar{x}, \bar{y}, \bar{u}, \bar{v})$ which solves the dual programs (2.1)-(2.2) with equal extrema. Any such solution $(\bar{x}, \bar{y}, \bar{u}, \bar{v})$ is bounded as follows:

$$\|\bar{x}, \bar{u}\|_1 \leq \left(\sum_{j=1}^p \lambda^j (x^j v^j + u^j y^j) \right) / \min_{\substack{1 \leq i \leq n \\ 1 \leq \ell \leq m}} \{\hat{v}_i, \hat{y}_\ell\}$$

Note that the requirement $u^j \geq 0$ in Theorem 2.3 is made because the monotonicity of F of Lemma 2.1 is established only on $R^n \times R_+^m$ and not on $R^n \times R^m$.

We give now a simple example illustrating the bounds of Theorem 2.2. This example shows that by a judicious parametrization of the point satisfying the required constraint qualification, tight bounds may be possible.

2.4 Example $\min x_1 + x_2 \quad \text{s.t. } y = x_2 - e^{x_1} \geq 0, x_1, x_2 \geq 0$

The dual problem is

$$\begin{aligned} \max \quad & x_1 + x_2 - u(x_2 - e^{x_1}) - vx \\ \text{s.t.} \quad & v_1 = 1 + ue^{x_1} \geq 0 \\ & v_2 = 1 - u \geq 0 \\ & u \geq 0 \end{aligned}$$

The primal-dual solution is $\bar{x}_1 = 0, \bar{x}_2 = 1, \bar{y} = 0, \bar{u} = 1, \bar{v}_1 = 2, \bar{v}_2 = 0$.

(a) To get a bound on $\|\bar{x}\|_1$ we need $v > 0$, so take $u = 0$ and hence

$$v_1 = v_2 = 1. \quad \text{Take } x_1 = \alpha \geq 0, x_2 = e^\alpha + \beta, \beta \geq 0 \text{ and hence}$$

$$y = \beta, xv + uy = \alpha + e^\alpha + \beta \text{ and}$$

$$1 = \|\bar{x}\|_1 \leq \inf_{\alpha \geq 0, \beta \geq 0} \{\alpha + e^\alpha + \beta\} = 1$$

(b) To get a bound on $\|\bar{y}\|_1$, take $x_1 = \alpha \geq 0$, $x_2 = e^\alpha$ and $u = 1$. Hence $y = 0$, $v_1 = 1 + e^\alpha$, $v_2 = 0$, $xv + uy = \alpha(1 + e^\alpha)$ and

$$0 = \|\bar{y}\|_1 \leq \inf_{\alpha \geq 0} \alpha(1 + e^\alpha) = 0$$

(c) To get a bound on $\|\bar{u}\|_1$ we need $y > 0$. So take $x_1 = 1 > 0$, $x_2 = \alpha + e$, $\alpha > 0$ and $u = \gamma \geq 0$. Hence $y = \alpha$, $v_1 = 1 + \gamma e$, $v_2 = 1 - \gamma \geq 0$, $xv + uy = 1 + e + \alpha$ and

$$1 = \|\bar{u}\|_1 \leq \inf_{\alpha > 0} \frac{1 + e + \alpha}{\alpha} = 1$$

(d) To get a bound on $\|\bar{v}\|_1$, take $x_1 = \alpha > 0$, $x_2 = e^\alpha$ and $u = 1$. Hence $y = 0$, $v_1 = 1 + e^\alpha$, $v_2 = 0$, $xv + uy = \alpha(1 + e^\alpha)$, and

$$2 = \|\bar{v}\|_1 \leq \inf_{\alpha > 0} \frac{\alpha(1 + e^\alpha)}{\alpha} = 2$$

We conclude by remarking that extensions of the results in this paper can also be established for the more general case in which the continuous monotone function F is replaced by a maximal monotone multifunction. Such extensions allow us to handle problem (2.1) with f and g nondifferentiable, convex and possibly taking the value of $+\infty$. Further extensions can also be proved in which R^k is replaced, for example, by any reflexive Banach space and R_+^k is replaced by a closed convex cone satisfying certain interiority/linearity properties [9, Corollary 1A]. We note also that the distributed constraint qualification of Theorems 1.2(a) and 1.3 which is sufficient for the boundedness result, can also be shown to be necessary in the linear case [7, Theorem 2.2] and even in the more general multivalued monotone case [9, Theorem 4].

References

1. L. E. J. Brouwer: "Über Abbildung von Mannigfaltigkeiten", *Mathematische Annalen* 71, 1910, 97-115.
2. R. W. Cottle: "Nonlinear programs with positively bounded Jacobians", *SIAM Journal of Applied Mathematics* 14, 1966, 147-158.
3. G. B. Dantzig, E. Eisenberg & R. W. Cottle: "Symmetric dual nonlinear programs", *Pacific Journal of Mathematics* 15, 1965, 809-812.
4. B. C. Eaves: "On the basic theorem of complementarity", *Mathematical Programming* 1, 1971, 68-75.
5. S. Karamardian: "The nonlinear complementarity problem with applications, part 2", *Journal of Optimization Theory and Applications* 4, 1969, 167-181.
6. O. L. Mangasarian: "Nonlinear programming", McGraw-Hill, New York, 1969.
7. O. L. Mangasarian: "Simple computable bounds for solutions of linear complementarity problems and linear programs", Technical Report #519, Computer Sciences Department, University of Wisconsin-Madison, October 1983, *Mathematical Programming Study*, to appear.
8. L. McLinden: "The complementarity problem for maximal monotone multifunctions", Chapter 17, 251-270 in "Variational inequalities and complementarity problems", R. W. Cottle, F. Giannessi & J.-L. Lions (editors), Wiley, New York, 1980.
9. L. McLinden; "Stable monotone variational inequalities", Technical Summary Report #2734, Mathematics Research Center, University of Wisconsin-Madison, August 1984.
10. N. Megiddo: "A monotone complementarity problem with feasible solutions but no complementary solutions", *Mathematical Programming* 12, 1977, 131-132.
11. J. J. Moré: "Coercivity conditions in nonlinear complementarity problems", *SIAM Review* 16, 1974, 1-16.
12. J. J. Moré: "Classes of functions and feasibility conditions in nonlinear complementarity problems", *Mathematical Programming* 6, 1974, 327-338.
13. R. T. Rockafellar: "A general correspondence between dual minimax problems and convex programs", *Pacific Journal of Mathematics* 25, 1968, 597-611.
14. D. R. Smart: "Fixed point theorems", Cambridge University Press, Cambridge, 1974.