

**CENTER FOR
PARALLEL OPTIMIZATION**

**ERROR BOUNDS AND STRONG UPPER
SEMICONTINUITY FOR MONOTONE
AFFINE VARIATIONAL INEQUALITIES**

by

M. C. Ferris & O. L. Mangasarian

Computer Sciences Technical Report #1056

November 1991

Error Bounds and Strong Upper Semicontinuity for Monotone Affine Variational Inequalities*

M.C. Ferris & O.L. Mangasarian

Computer Sciences Department

University of Wisconsin

1210 West Dayton Street

Madison, Wisconsin 53706

email: *ferris@cs.wisc.edu*, *olvi@cs.wisc.edu*

Technical Report # 1056

November 1991

Abstract. Global error bounds for possibly degenerate or nondegenerate monotone affine variational inequality problems are given. The error bounds are on an arbitrary point and are in terms of the distance between the given point and a solution to a convex quadratic program. For the monotone linear complementarity problem the convex program is that of minimizing a quadratic function on the nonnegative orthant. These bounds may form the basis of an iterative quadratic programming procedure for solving affine variational inequality problems. A strong upper semicontinuity result is also obtained which may be useful for finitely terminating any convergent algorithm by periodically solving a linear program.

Key words. Error bounds, upper semicontinuity, variational inequalities, linear complementarity problem

Abbreviated title. Error bounds for variational inequalities

*This material is based on research supported by Air Force Office of Scientific Research Grant AFOSR-89-0410 and National Science Foundation Grants CCR-9101801 and CCR-9157632

1 Introduction

We consider here the monotone affine variational inequality problem [Aus76, CGL80] of finding an \bar{x} in X such that

$$(x - \bar{x})^T(M\bar{x} + q) \geq 0, \forall x \in X := \{x \mid Ax \geq b, x \geq 0\} \quad (1)$$

where M is an $n \times n$ real positive semidefinite matrix (not necessarily symmetric) and A is an $m \times n$ real matrix. When X is the nonnegative orthant, the problem becomes the classical monotone linear complementarity problem (LCP) [CD68, Mur88]. Our principal concern here is: Given an arbitrary point x in \mathbb{R}^n , how far is it from the closed convex solution set \bar{X} of (1), assuming that \bar{X} is nonempty? Global error bounds for (1) have been given by [Pan86, Lemma 2] for positive definite M in terms of a “gradient projection” residual as well as by Tseng and Luo [LT92, Theorem 2.1] locally for a general monotone affine variational inequality. In [Man90b], global error bounds were also given for the case of positive semidefinite M in terms of a gradient projection residual multiplied by a term that involves the norm of the point x . In this work the error bound is motivated by the following formulation of (1) of finding an $\bar{x} \in X$ such that

$$(x - \bar{x})^T(Mx + q) \geq 0, \forall x \in X = \{x \mid Ax \geq b, x \geq 0\} \quad (2)$$

This is equivalent to (1) when M is positive semidefinite. Thus, given any point x in \mathbb{R}^n , we consider the convex quadratic program of finding a $y(x)$ such that

$$y(x) \in \arg \min_{y \in X} (y - x)^T(My + q) + \frac{\gamma}{2} \|y - x\|^2 \quad (3)$$

where $\gamma \geq 0$. Note that when $\gamma > 0$, problem (3) is a strongly convex quadratic program and hence has a unique solution $y(x)$. Furthermore, when $x \in \bar{X}$, it follows from the fact that M is positive semidefinite that

$$(y - x)^T(My + q) \geq (y - x)^T(Mx + q) \geq 0$$

Hence, the minimum value of zero can be achieved by setting $y = x$. Thus when $x \in \bar{X}$, and $\gamma > 0$, the unique solution of (3) is $y(x) = x$. The error bounds we propose here are based on the quantity $y(x) - x$ which is zero when $\bar{x} \in \bar{X}$ and $y(\bar{x}) = \bar{x}$. This prompts us to propose the following iterative procedure for solving (1)

$$x^{i+1} \in \arg \min_{x \in X} (x - x^i)^T(Mx + q) + \frac{\gamma}{2} \|x - x^i\|^2 \quad (4)$$

for which we have no convergence results at the present time. One advantage of (4) over the original problem (1) is the induced symmetry of the quadratic term which permits the potential use of successive overrelaxation (SOR) methods such as those of [DLM88] especially for the monotone linear complementarity problem, that is when X is the nonnegative orthant. In addition, (4) is strongly convex when $\gamma > 0$.

Our principal tool in deriving our bounds here is the error bound that was obtained in [MS86, Theorem 2.7] for the monotone linear complementarity problem. In Section 2 of this paper we give a simplified version of this error bound, and in Section 3 we derive our error bounds for (1), both for a possibly degenerate problem (Theorem 3) and a nondegenerate problem (Theorem 6). We also establish (Theorem 7) a strong upper semicontinuity result for the nondegenerate monotone affine variational inequality problem in terms of solutions of linear programs ((34) below) obtained by linearizing around any sequence $\{x^i\}$ which converges to a solution of the problem. From a certain \bar{i} onward, all solutions of the linear programs yield a solution of the variational inequality problem. This may be a useful result for finitely terminating any convergent algorithm for variational inequalities.

A word about our notation. For a vector x in the n -dimensional space \mathbb{R}^n , x_+ will denote the orthogonal projection on the nonnegative orthant \mathbb{R}_+^n , that is $(x_+)_i := \max\{x_i, 0\}$, $i = 1, \dots, n$. The norm $\|\cdot\|$ will denote the Euclidean norm, while other norms will be appropriately subscripted. The transpose of a matrix M will be denoted by M^T . For an $m \times n$ real matrix A , A_i will denote the i th row, while A_J will denote the set of rows A_i for $i \in J \subset \{1, \dots, m\}$. Similar notation, b_i and b_J is used for a vector b in \mathbb{R}^m . The identity matrix of arbitrary dimension will be denoted by I .

2 Preliminary Background

We begin by giving a simplification of the error bound of [MS86, Theorem 2.7] for the monotone linear complementarity problem

$$Nz + p \geq 0, z \geq 0, z^T(Nz + p) = 0 \quad (5)$$

where N is a $k \times k$ real positive semidefinite matrix, not necessarily symmetric. We shall assume that the solution set of (5), $SOL(N, p)$, is nonempty. By slight modification of the proofs of Lemma 2.5 and Theorem 2.7 of [MS86] and making use of Theorem 2.6 of [Man90a] we obtain the following simplified error bounds in terms of another “natural” residual $s(z)$ (see (8) below).

Theorem 1 (*Error bound for monotone linear complementarity problems*) *Let N be positive semidefinite, let the solution set $SOL(N, p)$ of (5) contain \bar{z} and let*

$$\hat{N} := \frac{1}{2}(N + N^T) \text{ and } d := 2\hat{N}\bar{z} + p \quad (6)$$

For any point z in \mathbb{R}^k there exists a $\bar{z}(z)$ in $SOL(N, p)$ such that [MS86, Theorem 2.7]

$$\|z - \bar{z}(z)\| \leq \sigma(N, p) \left(s(z) + s(z)^{\frac{1}{2}} \right) \quad (7)$$

where $s(z)$ is the residual

$$s(z) := \left\| \begin{pmatrix} -Nz - p, -z, z^T(Nz + p) \end{pmatrix}_+ \right\|, \quad (8)$$

and

$$\sigma(N, p) := \sqrt{k\nu(N, p)} \tau(N, p) \quad (9)$$

$$\nu(N, p) := \min_{\bar{z} \in \text{SOL}(N, p)} \|1, \bar{z}, N\bar{z} + p\| \quad (10)$$

$$\tau(N, p) := \sup_{(u, v, \xi, z)} \left\{ \|u, v, \xi, z\| \left| \begin{array}{l} \|N^T u + v - d\xi + \hat{N}^{\frac{1}{2}} z\|_1 = 1 \\ (u, v, \xi) \geq 0 \\ \text{Columns of } [N^T \ I \ d \ \hat{N}^{\frac{1}{2}}], \\ \text{corresponding to nonzero} \\ \text{elements of } (u, v, z, \xi), \\ \text{are linearly independent} \end{array} \right. \right\} \quad (11)$$

If the LCP (5) has some nondegenerate solution \hat{z} , that is $\hat{z} + N\hat{z} + p > 0$, then the bound (7) simplifies to [Man90a, Theorem 2.6]

$$\|z - \bar{z}(z)\| \leq \sigma(N, p)s(z), \quad (12)$$

with the term $\nu(N, p)$ deleted from (9), and the terms $\hat{N}^{\frac{1}{2}} z$ and $\hat{N}^{\frac{1}{2}}$ deleted from (11).

It was noted in [MS86, Example 2.9] that the square root term in the bound (7) is essential and cannot be dispensed with even locally.

In order to apply the above bounds to the monotone affine variational inequality problem (1) we state an equivalent characterization of (1) as a linear complementarity problem in \mathbb{R}^{n+m} as follows.

Proposition 2 *A point \bar{x} in \mathbb{R}^n solves the affine variational inequality (1) if and only if \bar{x} and some \bar{u} in \mathbb{R}^m solve the following linear complementarity problem*

$$w = \begin{bmatrix} M & -A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} x \\ u \end{bmatrix} + \begin{bmatrix} q \\ -b \end{bmatrix} \geq 0, \begin{bmatrix} x \\ u \end{bmatrix} \geq 0, w^T \begin{bmatrix} x \\ u \end{bmatrix} = 0 \quad (13)$$

Proof An \bar{x} is a solution of (1) if and only if $\bar{x} \in \arg \min_{x \in X} x^T(M\bar{x} + q)$. This is so if and only if \bar{x} and some \bar{u} satisfy the Karush-Kuhn-Tucker conditions (13) for the linear program $\min_{x \in X} x^T(M\bar{x} + q)$. \square

3 Error Bounds for the Monotone Affine Variational Inequality

We first derive the global error bounds (14) and (32) below, for the monotone variational inequality (1), by using the error bounds (7) and (12) on the equivalent LCP formulation (13) of (1).

Theorem 3 *(Global error bound for monotone affine variational inequalities) Let M be positive semidefinite, $\gamma \geq 0$, let (1) be solvable and let x be any given point in \mathbb{R}^n . Suppose one of the following three conditions holds:*

- (i) $\gamma > 0$, or
- (ii) $Mx - A^T u + q \geq 0$ for some $u \geq 0$, or
- (iii)

$$\left. \begin{array}{l} Ar \geq 0 \\ r \geq 0 \\ (M + M^T)r = 0 \end{array} \right\} \implies r = 0$$

Then there exists a solution $\bar{x}(x)$ of (1) such that

$$\|x - \bar{x}(x)\| \leq \tau(M, A, q, b)(t(x) + t(x)^{\frac{1}{2}}) + \|y(x) - x\| \quad (14)$$

where

$$y(x) \in \arg \min_{y \in X} (y - x)^T (My + q) + \frac{\gamma}{2} \|y - x\|^2, \quad (15)$$

$$t(x) := \left\| \left([M + \gamma I]^T (y(x) - x), -y(x)^T [M + \gamma I]^T (y(x) - x) \right)_+ \right\|, \quad (16)$$

$$\tau(M, A, q, b) := \sigma(N, p) \quad (17)$$

and σ is defined in (9) with

$$N = \begin{bmatrix} M & -A^T \\ A & 0 \end{bmatrix}, p = \begin{bmatrix} q \\ -b \end{bmatrix}, k = n + m \quad (18)$$

Remark Condition (iii) of the theorem is equivalent to every eigenvector r of $M + M^T$ corresponding to a zero eigenvalue which satisfies $Ar \geq 0$ having at least one negative component.

Proof Note first that (15) has a solution. This is trivial under assumption (i), by strong convexity and $X \neq \emptyset$. For the case when $\gamma = 0$, this follows from the facts that X is nonempty and the dual quadratic program [Man69], given by

$$\max_{y, u} \left\{ -y^T My - q^T x + b^T u \mid (M + M^T)y - M^T x - A^T u + q \geq 0, u \geq 0 \right\} \quad (19)$$

is feasible and hence the convex quadratic objective of (15) is bounded from below. That the dual problem (19) is feasible follows from assumption (ii) by taking $y = x$ in (19), or from assumption (iii) since this implies

$$(M + M^T)r = 0, Ar \geq 0, 0 \neq r \geq 0 \text{ has no solution } r$$

and hence by Tucker's Theorem [Man69, Theorem 2.4.3] there exist (y, u) such that

$$(M + M^T)y - A^T u > 0, u \geq 0$$

By multiplying this (y, u) by sufficiently large $\lambda > 0$ it follows that

$$(M + M^T)\lambda y - Mx - A^T\lambda u + q > 0, \lambda u \geq 0$$

and hence $(\lambda y, \lambda u)$ is feasible for (19).

It now follows from duality [Man69, Theorem 8.2.4] that if $y(x)$ is a solution of the convex quadratic program (15), then $y(x)$ and some $u(x)$ must also solve the dual quadratic program

$$\begin{aligned} & \underset{y, u}{\text{maximize}} && -y^T M y + \frac{\gamma}{2} \|y - x\|^2 - \gamma y^T (y - x) - q^T x + b^T u \\ & \text{subject to} && (M + M^T)y + \gamma(y - x) - M^T x - A^T u + q \geq 0, u \geq 0 \end{aligned} \quad (20)$$

and

$$(y(x) - x)^T (M y(x) + q) = -y(x)^T M y(x) - \gamma y(x)^T (y(x) - x) - q^T x + b^T u(x) \quad (21)$$

We now apply the error bound, Theorem 1, to the linear complementarity problem (13) (which is equivalent to (1)) at the point $z = \begin{bmatrix} y(x) \\ u(x) \end{bmatrix}$ with N and p as defined in (18). We note first that $s(z)$ as defined by (8) is

$$s(z) = \left\| \left(\begin{bmatrix} -M y(x) + A^T u(x) - q \\ -A y(x) + b \end{bmatrix}, \begin{bmatrix} -y(x) \\ -u(x) \end{bmatrix}, y(x)^T M y(x) + q^T y(x) - b^T u(x) \right)_+ \right\| \quad (22)$$

From the equality of (21) we have that

$$y(x)^T M y(x) + q^T y(x) - b^T u(x) = -(y(x) - x)^T [M + \gamma I] y(x) \quad (23)$$

From the feasibility of $y(x)$, $u(x)$ for (20) we get

$$-M y(x) + A^T u(x) - q \leq [M + \gamma I]^T (y(x) - x) \quad (24)$$

Using (23), (24), $y(x) \geq 0$, $u(x) \geq 0$ and $A y(x) \geq b$ in (22) we obtain (upon invoking the monotonicity of the Euclidean norm)

$$s(z) \leq \left\| \left([M + \gamma I]^T (y(x) - x), -y(x)^T [M + \gamma I]^T (y(x) - x) \right)_+ \right\| = t(x) \quad (25)$$

where the last equality follows from the definition (16) of $t(x)$. Using the expression (25) in (7) we obtain that there exists

$$\begin{bmatrix} \bar{x}(x) \\ \bar{u}(x) \end{bmatrix} := \begin{bmatrix} \bar{x}(y(x), u(x)) \\ \bar{u}(y(x), u(x)) \end{bmatrix}$$

which solves (13) such that

$$\|y(x) - \bar{x}(x)\| \leq \left\| \begin{bmatrix} y(x) - \bar{x}(x) \\ u(x) - \bar{u}(x) \end{bmatrix} \right\| \leq \tau(M, A, q, b)(t(x) + t(x)^{\frac{1}{2}}) \quad (26)$$

However, since

$$\|x - \bar{x}(x)\| \leq \|x - y(x)\| + \|y(x) - \bar{x}(x)\| \quad (27)$$

it follows that

$$\|x - \bar{x}(x)\| \leq \tau(M, A, q, b)(t(x) + t(x)^{\frac{1}{2}}) + \|x - y(x)\|$$

which gives the required result upon noting that $\bar{x}(x)$ solves (1). \square

We point out the difference between the two cases $\gamma > 0$ and $\gamma = 0$. In the former case, the error bound (14) would seem to be weaker than the corresponding one for $\gamma = 0$. However, if $x \in \bar{X}$, it is easy to see that $y(x) = x$ when $\gamma > 0$, so the corresponding error bound is zero on the solution set. There is no such guarantee for the case $\gamma = 0$, since there may be other solutions to (15). The following lemma and example make this clearer.

Lemma 4 *Suppose that M is positive semidefinite and \bar{x} is in the solution set \bar{X} of (1). Let*

$$\bar{Y} := \left\{ x \in X \mid \exists u \geq 0, Mx + q - A^T u \geq 0, (M + M^T)(x - \bar{x}) = 0, (M\bar{x} + q)^T(x - \bar{x}) = 0 \right\},$$

$$\bar{Z} := \arg \min_{y \in X} (y - \bar{x})^T (My + q)$$

and

$$\bar{N} := \{x \mid M(x - \bar{x}) = 0\}$$

Then $\bar{X} \subset \bar{Y} \subset \bar{Z}$, $\bar{Z} \cap \bar{N} \subset \bar{X}$ and hence

$$\bar{X} \cap \bar{N} = \bar{Y} \cap \bar{N} = \bar{Z} \cap \bar{N}$$

Proof In order to show that $\bar{X} \subset \bar{Y}$, suppose $\bar{y} \in \bar{X}$. Then

$$(x - \bar{y})^T (M\bar{y} + q) \geq 0, \forall x \in X$$

The fact that M is positive semidefinite and $\bar{x} \in X$ gives

$$(\bar{x} - \bar{y})^T (M\bar{x} + q) \geq (\bar{x} - \bar{y})^T (M\bar{y} + q) \geq 0$$

Since $\bar{x} \in \bar{X}$, it follows that $(x - \bar{x})^T (M\bar{x} + q) \geq 0$ for all $x \in X$, so $\bar{y} \in X$ implies $(\bar{y} - \bar{x})^T (M\bar{x} + q) \geq 0$ and hence

$$(\bar{y} - \bar{x})^T (M\bar{x} + q) = 0 \quad (28)$$

Furthermore,

$$\begin{aligned} 0 &\leq (\bar{y} - \bar{x})^T M(\bar{y} - \bar{x}) \\ &= (\bar{y} - \bar{x})^T (M\bar{y} + q) - (\bar{y} - \bar{x})^T (M\bar{x} + q) \\ &= (\bar{y} - \bar{x})^T (M\bar{y} + q) \\ &\leq 0 \end{aligned}$$

the second equality following from (28) and the final inequality following from the fact that $\bar{y} \in \bar{X}$. Hence, $(\bar{y} - \bar{x})^T M(\bar{y} - \bar{x}) = 0$, giving

$$(M + M^T)(\bar{y} - \bar{x}) = 0$$

The proof that $\bar{X} \subset \bar{Y}$ is complete if we can exhibit a $u \geq 0$ such that $M\bar{y} + q \geq A^T u$. However, by Proposition 2, $\bar{y} \in \bar{X}$ implies that \bar{y} and some \bar{u} solve (13), which by feasibility of \bar{y} and \bar{u} shows

$$M\bar{y} - A^T \bar{u} + q \geq 0, \bar{u} \geq 0$$

as required.

We now show $\bar{Y} \subset \bar{Z}$. It is easy to see that $\bar{x} \in \bar{Z}$ and so it follows from the characterization of the solution set of a convex quadratic program given in [Man88, Corollary 1] that

$$\bar{Z} = \left\{ x \in X \mid (M + M^T)(x - \bar{x}) = 0, (M\bar{x} + q)^T(x - \bar{x}) = 0 \right\} \quad (29)$$

The inclusion $\bar{Y} \subset \bar{Z}$ is now obvious.

To establish the last conclusions of the lemma, let $x \in \bar{Z}$. By the definition of \bar{Z} , x and some u must satisfy the following Karush-Kuhn-Tucker optimality conditions

$$\begin{aligned} (M + M^T)x + q - M^T \bar{x} - A^T u &\geq 0 & Ax - b &\geq 0 \\ x^T((M + M^T)x + q - M^T \bar{x} - A^T u) &= 0 & u^T(Ax - b) &= 0 \\ x &\geq 0 & u &\geq 0 \end{aligned} \quad (30)$$

Hence, if $x \in \bar{Z} \cap \bar{N}$, it follows from (29) that $M^T(x - \bar{x}) = 0$ and consequently (30) becomes (13) and so x solves (1), that is $x \in \bar{X}$. This gives the desired inclusion $\bar{Z} \cap \bar{N} \subset \bar{X}$. Since we already have that $\bar{X} \subset \bar{Y} \subset \bar{Z}$, it follows that

$$\bar{X} \cap \bar{N} = \bar{Y} \cap \bar{N} = \bar{Z} \cap \bar{N}$$

□

The following example from [MS86] shows that the inclusions in Lemma 4 may be strict, even for the case when the linear complementarity problem (13) associated with (1) has a nondegenerate solution.

Example 5

$$M = \begin{bmatrix} 0 & -1 \\ 1 & 1 \end{bmatrix}, q = \begin{bmatrix} 1 \\ -2 \end{bmatrix}, X = \mathbb{R}_+^2$$

Then $\bar{X} = \{(1, 1)\}$, $\bar{Y} = \{(x, 1) \mid x \geq 1\}$ and $\bar{Z} = \{(x, 1) \mid x \geq 0\}$.

If the monotone affine variational inequality (1) is nondegenerate in the sense that the corresponding LCP (13) has some nondegenerate solution, then by using the corresponding error bound (12) of Theorem (1), the error bound (14) given above simplifies as follows.

Theorem 6 (*Error bound for nondegenerate monotone affine variational inequality*) Let M be positive semidefinite, $\gamma \geq 0$ and let (1) be solvable with \hat{x} being some nondegenerate solution of (1), that is \hat{x} and a corresponding \hat{u} solve (13) such that

$$\hat{x} + M\hat{x} - A^T\hat{u} + q > 0, \hat{u} + A\hat{x} - b > 0 \quad (31)$$

Suppose one of the following three conditions holds:

- (i) $\gamma > 0$, or
- (ii) $Mx - A^T u + q \geq 0$ for some $u \geq 0$, or
- (iii)

$$\left. \begin{array}{l} Ar \geq 0 \\ r \geq 0 \\ (M + M^T)r = 0 \end{array} \right\} \implies r = 0$$

Then there exists a solution $\bar{x}(x)$ of (1) such that

$$\|x - \bar{x}(x)\| \leq \tau(M, A, q, b)t(x) + \|y(x) - x\| \quad (32)$$

where

$$y(x) \in \arg \min_{y \in X} (y - x)^T (My + q) + \frac{\gamma}{2} \|y - x\|^2,$$

$$t(x) := \left\| \left([M + \gamma I]^T (y(x) - x), -y(x)^T [M + \gamma I]^T (y(x) - x) \right)_+ \right\|,$$

and

$$\tau(M, A, q, b) := \sqrt{k} \sup_{(u, v, \xi)} \left\{ \|u, v, \xi\| \left| \begin{array}{l} \|N^T u + v - d\xi\|_1 = 1 \\ (u, v, \xi) \geq 0 \\ \text{Columns of } [N^T \ I \ d], \\ \text{corresponding to nonzero} \\ \text{elements of } (u, v, \xi), \\ \text{are linearly independent} \end{array} \right. \right\}$$

with N and p and k defined by (18).

Remark Theorems 3 and 6 can be specialized to the monotone linear complementarity problem by taking X to be the nonnegative orthant in \mathbb{R}^n . In this case, $y(x)$ as defined by (3) simplifies to

$$y(x) \in \arg \min_{y \geq 0} (y - x)^T (My + q) + \frac{\gamma}{2} \|y - x\|^2$$

We now derive a strong upper semicontinuity result for the nondegenerate monotone affine variational inequality problem based on a similar result for the monotone linear complementarity problem [FM89].

Theorem 7 (*Strong upper semicontinuity of nondegenerate monotone affine variational inequalities*) Let M be positive semidefinite and let (1) be solvable with \hat{x} being **some** nondegenerate solution of (1), that is \hat{x} and a corresponding \hat{u} solve (13) and such that (31) holds. Let $\{x^i\}$ be a sequence in X converging to an $\bar{x} \in \bar{X}$ such that there exist $\{u^i\}$ satisfying

$$Mx^i - A^T u^i + q \geq 0, u^i \geq 0 \quad (33)$$

Then for sufficiently large i , $y^i \in \bar{X}$, where (y^i, u^i) solve

$$\min_{y,u} \left\{ ((M + M^T)x^i + q)^T y - b^T u \mid My - A^T u + q \geq 0, Ay \geq b, y, u \geq 0 \right\} \quad (34)$$

Proof By Corollary 14 of [FM89], there exists an \bar{i} , such that for all $i \geq \bar{i}$

$$\arg \min_{y,u} \left\{ ((M + M^T)x^i + q)^T y - b^T u \mid My - A^T u + q \geq 0, Ay \geq b, y, u \geq 0 \right\} \subset SOL(N, p)$$

where $SOL(N, p)$ is the solution set of (13). That is any solution of a linearization of the quadratic program

$$\min_{y,u} \left\{ y^T My + q^T y - b^T u \mid My - A^T u + q \geq 0, Ay \geq b, y, u \geq 0 \right\} \quad (35)$$

associated with the LCP (13) around (x^i, u^i) yields an exact solution of (13) and hence x^i is an exact solution of (1). \square

Remark It is interesting to note that since the quadratic program (35) is linear in u , the sequence $\{u^i\}$ is not needed in solving the linearization (34). From a practical point of view, this leads to finite termination of any convergent algorithm for solving (1) in the nondegenerate case by periodically solving the linear program of (34).

4 Conclusion

We have presented error bounds for the monotone affine variational inequality problem, both for the degenerate and nondegenerate cases. These bounds may potentially be useful in constructing an iterative quadratic programming algorithm for solving affine variational inequalities. We have also derived a strong upper semicontinuity results for the nondegenerate monotone affine variational inequality problem which may be useful for finitely terminating any convergent algorithm by periodically solving a linear program.

References

- [Aus76] A. Auslender. *Optimization Méthodes Numériques*. Masson, Paris, 1976.
- [CD68] R.W. Cottle and G. Dantzig. Complementary pivot theory of mathematical programming. *Linear Algebra and its Applications*, 1:103–125, 1968.

- [CGL80] R.W. Cottle, F. Giannessi, and J.-L. Lions. *Variational Inequalities and Complementarity Problems: Theory and Applications*. John Wiley & Sons, New York, 1980.
- [DLM88] R. De Leone and O.L. Mangasarian. Serial and parallel solution of large scale linear programs by augmented Lagrangian successive overrelaxation. In A. Kurzbaniski, K. Neumann, and D. Pallaschke, editors, *Optimization, Parallel Processing and Applications*, pages 103–124, Berlin, 1988. Springer-Verlag. Lecture Notes in Economics and Mathematical Systems 304.
- [FM89] M.C. Ferris and O.L. Mangasarian. Minimum principle sufficiency. Technical Report 853, Computer Sciences Department, University of Wisconsin, Madison, Wisconsin 53706, June 1989. To appear in *Mathematical Programming*.
- [LT92] Z.-Q. Luo and P. Tseng. Error bound and convergence analysis of matrix splitting algorithms for the affine variational inequality problem. *SIAM Journal on Optimization*, 2(1), February 1992.
- [Man69] O.L. Mangasarian. *Nonlinear Programming*. McGraw-Hill, New York, 1969.
- [Man88] O.L. Mangasarian. A simple characterization of solution sets of convex programs. *Operations Research Letters*, 7(1):21–26, 1988.
- [Man90a] O.L. Mangasarian. Error bounds for nondegenerate monotone linear complementarity problems. *Mathematical Programming*, pages 437–445, 1990.
- [Man90b] O.L. Mangasarian. Global error bounds for monotone affine variational inequality problems. Technical Report 998, Computer Sciences Department, University of Wisconsin, Madison, Wisconsin 53706, January 1990. To appear in *Linear Algebra and Its Applications*.
- [MS86] O.L. Mangasarian and T.-H. Shiau. Error bounds for monotone linear complementarity problems. *Mathematical Programming*, 36:81–89, 1986.
- [Mur88] K.G. Murty. *Linear Complementarity, Linear and Nonlinear Programming*. Helderman-Verlag, Berlin, 1988.
- [Pan86] J.S. Pang. Inexact Newton methods for the nonlinear complementarity problem. *Mathematical Programming*, 36(1):54–71, 1986.